

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 48/00, C12N 15/63	A1	(11) International Publication Number: WO 97/06828
		(43) International Publication Date: 27 February 1997 (27.02.97)
(21) International Application Number: PCT/US96/13360 (22) International Filing Date: 16 August 1996 (16.08.96) (30) Priority Data: 60/002,490 18 August 1995 (18.08.95) US 60/002,479 18 August 1995 (18.08.95) US (71) Applicant (for all designated States except US): SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH [US/US]; 1275 York Avenue, New York, NY 10021 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROTHMAN, James, E. [US/US]; Apartment 10B, 402 East 64th Street, New York, NY 10021 (US). HARTL, F., Ulrich [DE/US]; 20 Quarry Lane, Irvington, NY 10533 (US). HOE, Mee, H. [MY/US]; 312 East 66th Street, New York, NY 10021 (US). HOUGHTON, Alan [US/US]; Apartment 7C, 402 East 64th Street, New York, NY 10021 (US). TAKEUCHI, Yoshizumi [JP/US]; Apartment 24M, 1635 York Avenue, New York, NY 10021 (US). MAYHEW, Mark [GB/US]; Apartment 16A, 10 River Road, New York, NY 10044 (US).		(74) Agents: LARSON, Marina, T. et al.; Oppedahl & Larson, Suite 309, 1992 Commerce Street, Yorktown Heights, NY 10598 (US). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: METHOD FOR TREATMENT OF CANCER AND INFECTIOUS DISEASES AND COMPOSITIONS USEFUL IN SAME		
(57) Abstract <p>Administration of expressible polynucleotides encoding eukaryotic heat shock proteins to mammalian cells leads to the stimulation of an immune response to antigens present in those cells. This makes it possible to stimulate an immune response to target antigens, including target tumor antigens or antigens associated with an infectious disease, without having to isolate a unique antigen or antigen-associated heat shock protein for each target antigen by administering to a mammalian subject or to a group of mammalian cells containing the antigen, an expressible polynucleotide encoding a heat shock protein. The expressed heat shock protein may have the same structure as native heat shock proteins, or may be a modified form adapted to control the trafficking of the expressed heat shock protein within the cells.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DESCRIPTIONMETHOD FOR TREATMENT OF CANCER AND INFECTIOUS DISEASES
AND COMPOSITIONS USEFUL IN SAME

The invention described herein was made in the course of work under NIH Core Grant No. CA 08748. The United States government may have certain rights in this invention.

5

BACKGROUND OF THE INVENTION

This application relates to the use of heat shock proteins and similar peptide-binding proteins to stimulate an immunological response against antigens, including cancer-related antigens, autoimmune antigens and infectious disease antigens, found in a mammalian host.

Heat shock proteins were originally observed to be expressed in increased amounts in mammalian cells which were exposed to sudden elevations of temperature, while the expression of most cellular proteins is significantly reduced. It has since been determined that such proteins are produced in response to various types of stress, including glucose deprivation. As used herein, the term "heat shock protein" will be used to encompass both proteins that are expressly labeled as such as well as other stress proteins, including homologs of such proteins that are expressed constitutively (i.e., in the absence of stressful conditions). Examples of heat shock proteins include BiP (also referred to as grp78), hsp/hsc70, gp96 (grp94), hsp60, hsp40 and hsp90.

Heat shock proteins have the ability to bind other proteins in their non-native states, and in particular to bind nascent proteins emerging from the ribosomes or extruded into endoplasmic reticulum. Hendrick and Hartl, *Ann. Rev. Biochem.* 62: 349-384 (1993); Hartl, *Nature* 381: 571-580 (1996). Further, heat shock proteins have been shown to play an important role in the proper folding and assembly of proteins in the cytosol, endoplasmic reticulum

- 2 -

and mitochondria; in view of this function, they are referred to as "molecular chaperones". Frydman et al., *Nature* 370: 111-117 (1994); Hendrick and Hartl., *Ann. Rev. Biochem.* 62:349-384 (1993); Hartl., *Nature* 381:571-580 (1996).

For example, the protein BiP, a member of a class of heat shock proteins referred to as the hsp70 family, has been found to bind to newly synthesized, unfolded μ immunoglobulin heavy chain prior to its assembly with light chain in the endoplasmic reticulum. Hendershot et al., *J. Cell Biol.* 104:761-767 (1987). Another heat shock protein, gp96, is a member of the hsp90 family of stress proteins which localize in the endoplasmic reticulum. Li and Srivastava, *EMBO J.* 12:3143-3151 (1993); Mazzarella and Green, *J. Biol. Chem.* 262:8875-8883 (1987). It has been proposed that gp96 may assist in the assembly of multi-subunit proteins in the endoplasmic reticulum. Wiech et al., *Nature* 358:169-170 (1992).

It has been observed that heat shock proteins prepared from tumors in experimental animals were able to induce immune responses in a tumor-specific manner; that is to say, heat shock protein purified from a particular tumor could induce an immune response in an experimental animal which would inhibit the growth of the same tumor, but not other tumors. Srivastava and Maki, 1991, *Curr. Topics Microbiol.* 167: 109-123 (1991). The source of the tumor-specific immunogenicity has not been confirmed. Genes encoding heat shock proteins have not been found to exhibit tumor-specific DNA polymorphism. Srivastava and Udon, *Curr. Opin. Immunol.* 6:728-732 (1994). High resolution gel electrophoresis has indicated that gp96 may be heterogeneous at the molecular level. Feldweg and Srivastava, *Int. J. Cancer* 63:310-314 (1995). Evidence suggests that the source of heterogeneity may be populations of small peptides adherent to the heat shock protein, which may number in the hundreds. *Id.* It has been proposed that a wide diversity of peptides adherent to tumor-synthesized heat shock proteins may render such proteins capable of eliciting an immune response in

- 3 -

subjects having diverse HLA phenotypes, in contrast to more traditional immunogens which may be somewhat HLA-restricted in their efficacy. *Id.*

5 Lukacs et al., *J. Exp. Med.* 178: 343-348 (1993),
have reported the transfection of tumor cells with a myco-
bacterial heat shock protein-encoding gene, and the obser-
vation that the transfected cells lose tumorigenicity and
induce what appears to be T-cell mediated protection against
tumors in mice immunized using the transfected cells.
10 Lukacs et al. suggest that the loss of tumorigenicity could
result from the interaction of the heat shock protein with
p53 via increased efficiency of chaperone activity to pro-
duce proper folding and conformation of otherwise ineffec-
tive p53 protein. They further suggest that the highly
15 immunogenic nature of the 65 kD bacterial hsp enhances the
recognition of other, tumor-associated antigen molecules.

It has been suggested in the literature that myco-
bacterial heat shock proteins may play a role in the onset
of autoimmune diseases such as rheumatoid arthritis. Thus,
20 the practical utility of such bacterial proteins in vaccines
for the treatment of humans is questionable. It is an
object of the present invention to provide vaccine composi-
tions which can be used to stimulate an immune response to
antigens, including tumor and infectious disease antigens,
25 present in mammalian cells without the introduction of
mycobacterial proteins.

SUMMARY OF THE INVENTION

30 It has now been found that administration of
expressible polynucleotides encoding eukaryotic heat shock
proteins to mammalian cells leads to the stimulation of an
immune response to antigens present in those cells. This
makes it possible to stimulate an immune response to treat a
subject's disease condition, including an immune response to
35 a tumor or an infectious disease, without having to isolate
or characterize an antigen associated with the disease.
Thus, the present invention provides a method for stimulat-
ing a therapeutic or prophylactic immune response in a

mammalian subject by treating the subject or a group of cells from the subject with an expressible polynucleotide encoding a eukaryotic heat shock protein. The expressed heat shock protein may have the same structure as native heat shock proteins, or may be a modified form adapted to control the trafficking of the expressed heat shock protein within the cells.

BRIEF DESCRIPTION OF THE DRAWINGS

10 Figs. 1A-H show tumor formation in mice injected with CMS-5 sarcoma cells and CMS-5 sarcoma cells acutely transfected with BiP-expressing vectors; and

15 Figs. 2A-F show tumor formation in mice injected with CMS-5 sarcoma cells and CMS-5 sarcoma cells stably transfected with BiP-expressing vectors.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method for inducing or enhancing an immune response to an antigen present in a target group of cells without requiring prior identification, characterization or isolation of the antigen. The method of the invention may be used to induce an immune response to the antigen in the case where there is no existing detectable disease-related immune response prior to practicing the method, or to enhance the disease-related immune response to a greater level. For purposes of simplicity, the specification and claims of this application will use the term "stimulating an immune response" to encompass both inducing a new immune response and enhancing a pre-existing immune response.

30 We have found that administration of expressible polynucleotides encoding eukaryotic heat shock proteins to mammalian cells leads to the stimulation of an immune response to antigens present in those cells. While we do not intend to be bound by any particular mechanism, the full potential of our observations is best understood in the context of two mechanistic models which explain the observations.

The first mechanism is based on the fact that heat shock proteins like BiP and gp96 are normally localized in the endoplasmic reticulum and are known to bind peptides and proteins in this location, and the belief that such binding may be a step in the presentation of an antigen to the immune system. Increasing the concentration of one or more heat shock proteins resident in the endoplasmic reticulum may cause an improvement in the efficiency of this process, and thus an increased immune response to the antigens present in the transfected cells.

The second mechanism is based upon the understanding that heat shock proteins and other proteins secreted from the endoplasmic reticulum may be recaptured by receptors known as erd-2 (KDEL) receptors and returned to the endoplasmic reticulum. This recapture process requires the presence of a specific signal on the heat shock protein such XDEL. XDEL is the single letter amino-acid code for the amino acid sequence variable-Asp-Glu-Leu. Specific carboxy-terminal signal sequences are KDEL (Lys-Asp-Glu-Leu) and HDEL (His-Asp-Glu-Leu). By eliminating this signal in heat shock proteins which bind to antigens or by otherwise interfering with the ability of the erd-2 receptor to recapture these heat shock proteins, the amount of antigen-associated heat shock protein secreted by a cell and therefore accessible to the immune system can be increased.

In accordance with either of these mechanisms, the stimulation of the immune response which is observed when an expressible polynucleotide encoding a heat shock protein is expressed in the cell flows from an increase in the amount of antigen-associated heat shock protein. Thus, in accordance with one embodiment of the invention there is provided a method for stimulating an immune response to an antigen present in a target group of mammalian cells, comprising administering to the target group of cells a composition effective to increase the amount of antigen-associated heat shock protein present in the cells. This can be accomplished in several ways.

First, the amount of antigen-associated heat shock protein can be increased by administering a polynucleotide encoding a heat shock protein which is expressed to produce recombinant heat shock protein that is processed and transported within the cell in the same manner as wild-type heat shock protein.

Second, the amount of antigen-associated heat shock protein can be increased by administering an expressible polynucleotide encoding a heat shock protein in a modified form adapted to control the trafficking of the expressed heat shock protein within the cells. This would include alterations of heat shock protein which either facilitate its secretion from the cell or its localization and/or retention on the cell. For example, secretion may be facilitated by mutating or eliminating portions of the heat shock protein that serve to retain the heat shock protein in the cell (for example by deleting sequences recognized by the erd-2 receptor such as KDEL or a functionally equivalent sequence or by adding an agent that interferes with binding of heat shock protein to erd-2), or by supplying mutated non-heat shock proteins (e.g. erd-2 mutants) that increase heat shock protein secretion. Alternatively localization and retention may be accomplished by increasing the number and/or strength of retention or retrieval signals (e.g. increasing erd-2 levels) or my by adding to a heat shock protein, a membrane anchor containing additional retention or retrieval sequence as well as transmembrane and cytoplasmic domains (e.g., KKXX at the C-terminal end, or XXRR at the N-terminal end or function equivalents thereof.)

Third, the amount of an antigen-associated heat shock protein can be increased by administering an expressible polynucleotide encoding a peptide or peptide which competitively inhibits the interaction of heat shock proteins with cellular elements to modify the trafficking of those heat shock proteins within the cell.

Preparation of Expressible Polynucleotides

The construction of expressible polynucleotides for use in the three embodiments of the invention share many common elements and use generally well known techniques.

5 These techniques are set forth in numerous literature references known to persons skilled in the art and so will not be repeated exhaustively here.

Expressible polynucleotide encoding a heat shock protein which is expressed to produce recombinant heat shock
10 protein that is processed and transported within the cell in the same manner as wild-type heat shock protein can be constructed by incorporating a cDNA encoding a wild-type heat shock protein or a modified heat shock protein that retains both its heat shock protein functionality and the
15 XDEL signal sequence into a mammalian expression vector. Genes for various mammalian heat shock proteins have been cloned and sequenced, including, but not limited to, gp96, human: Genebank Accession No. X15187; Maki et al., *Proc. Nat'l Acad. Sci.* 87: 5658-5562 (1990), mouse: Genebank
20 Accession No. M16370; Srivastava et al., *Proc. Nat'l Acad. Sci.* 84:3807-3811 (1987)); BiP, mouse: Genebank Accession No. U16277; Haas et al., *Proc. Nat'l Acad. Sci. U.S.A.* 85: 2250-2254 (1988), human: Genebank Accession No. M19645; Ting et al., *DNA* 7: 275-286 (1988); hsp70, mouse: Genebank
25 Accession No. M35021; Hunt et al., *Gene* 87: 199-204 (1990), human: Genebank Accession No. M24743; Hunt et al., *Proc. Nat'l Acad. Sci. U.S.A.* 82: 6455-6489 (1995); and hsp40 human: Genebank Accession No. D49547; Ohtsuka K., *Biochem Biophys. Res. Commun.* 197: 235-240 (1993).

30 In general, any type of mammalian expression vector can be used, although those with the highest transfection and expression efficiencies are preferred to maximize the levels of expression. Specific types of vectors which can be employed include herpes simplex viral
35 based vectors: pHSV1 (Geller et al. *Proc. Natl. Acad. Sci.* 87:8950-8954 (1990)); recombinant retroviral vectors: MFG (Jaffee et al. *Cancer Res.* 53:2221-2226-(1993)); Moloney-based retroviral vectors: LN, LNSX, LNCX, LXSX (Miller and

- 8 -

Rosman Biotechniques 7:980-989 (1989)); vaccinia viral vector: MVA (Sutter and Moss Proc. Natl. Acad. Sci. 89:10847-10851 (1992)); recombinant adenovirus vectors : pJM17 (Ali et al Gene Therapy 1:367-384 (1994)), (Berkner K. L. Biotechniques 6:616-624 1988); second generation adenovirus vector: DE1/DE4 adenoviral vectors (Wang and Finer Nature Medicine 2:714-716 (1996)); and Adeno-associated viral vectors: AAV/Neo (Muro-Cacho et al. J. Immunotherapy 11:231-237 (1992)). Specific suitable expression systems for this purpose include pCDNA3 (In-Vitrogen), plasmid AH5 (which contains the SV40 origin and the adenovirus major late promoter), pRC/CMV (In-Vitrogen), pCMU II (Paabo et al., EMBO J. 5: 1921-1927 (1986)), pZip-Neo SV (Cepko et al., Cell 37: 1053-1062 (1984)) and pSR α (DNAX, Palo Alto, CA).

Expressible polynucleotides encoding any of the various modified forms heat shock proteins described above are constructed using similar techniques. For example, heat shock proteins can be produced which are modified to delete or block the carboxy-terminal XDEL sequence. Such a heat shock protein is adapted to permit the heat shock protein to be secreted from the endoplasmic reticulum and to interfere with the return of the protein to the endoplasmic reticulum that is mediated by the erd-2 receptors because the signal peptide recognized by the erd-2 receptors is missing. Proteins of this type are referred to in the specification and claims of this application as XDEL-negative heat shock proteins.

Expressible XDEL-negative polynucleotides can be constructed by deleting the nucleotides encoding the amino acid sequence XDEL at the carboxy-terminus of the heat shock protein. This can be accomplished by PCR amplification of the wild-type cDNA using a primer that hybridizes in the region immediately adjacent to the portion coding for the XDEL signal, combined with a restriction site to be used for cloning the product into a desired vector for expression of the modified heat shock protein in mammalian cells as described above.

- 9 -

As an alternative to the use of expressible polynucleotides encoding deletion mutants of heat shock proteins, the method of the invention may also employ expressible polynucleotides which encode heat shock proteins in which the carboxy-terminal signal group XDEL is masked by the addition of additional amino acids to the carboxy-terminal end. The masking amino acids can serve only the masking function, or can be derived from an infectious agent (for example E7/E6 from human papilloma virus) in which case the masking amino acids may also function as an antigen.

Addition of masking amino acids can be accomplished as described in Munro et al., *Cell* 48: 899-907 (1987) in which a mutant (SAGGL) having two amino acids added after the XDEL was prepared by cloning in appropriate restriction sites to permit substitution of a sequence including the additional bases, or by PCR amplification of the heat shock protein using a primer that hybridizes with the bases encoding the XDEL signal and which has additional bases encoding the added nucleotides inserted between the XDEL-encoding bases and the stop codon. The number of additional amino acids does not matter, so long as the added amino acids do not alter the function of the heat shock protein. The nucleotides encoding the masked heat shock protein are then introduced into a mammalian expression vector as described above.

Another method which may be used to increase the amount of antigen-associated heat shock protein in the cells is to modify cytosolic heat shock proteins with an amino-terminal signal which causes the protein to be taken up into the endoplasmic reticulum where it can gain access to the secretory pathway. Amino-terminal signals which can accomplish this function are described in the art, including for example in von Heijne, G., *J. Mol. Biol.* 184: 99-105 (1985). These signals generally comprise a positively-charged amino acid or group of a few amino acids at the amino terminal end of the peptide, a hydrophobic core region and a third region of greater polarity than the hydrophobic region. Such additional signal sequences can be introduced using PCR with

- 10 -

a modified primer that includes the bases for the desired amino-terminal amino acids. A representative cytosolic heat shock protein which might be modified in this manner is hsp 70.

5 The expressible polynucleotide used in the method of the present invention may also encode a peptide or protein that competitively inhibits the interaction of heat shock proteins with cellular elements to modify the trafficking of those heat shock proteins within the cell. In particular, the expressible polynucleotide may encode a peptide or protein that includes a carboxy-terminal XDEL signal which will compete with heat shock proteins, including endogenous heat shock proteins, for binding to the erd-2 receptors. This competitive peptide may itself be a heat shock protein, for example BiP, that retains the XDEL signal sequence, or it may be some other peptide or protein with a carboxy-terminal XDEL sequence.

Administration of Expressible Polynucleotides

20 The resulting expressible polynucleotide is delivered into cells of the subject by *ex vivo* or *in vivo* methods, for example as part of a viral vector as described above, or as naked DNA. Suitable methods include injection directly into tissue and tumors, transfection using liposomes (Fraley et al., *Nature* 370: 111-117 (1980)), receptor-mediated endocytosis (Zatloukal, et al., *Ann. N.Y. Acad. Sci.* 660: 136-153 (1992)), particle bombardment-mediated gene transfer (Eisenbraun et al., *DNA & Cell. Biol.* 12: 791-797 (1993)) and transfection using peptide presenting bacteriophage. Barry et al. *Nature Medicine* 2: 299-305 (1996).

35 The expressible polynucleotide may be administered to a subject in need of treatment in order to obtain a therapeutic immune response. The term "therapeutic immune response", as used herein, refers to an increase in humoral and/or cellular immunity, as measured by standard techniques, which is directed toward an antigen associated with the subject's disease condition. Preferably, but not by

- 11 -

way of limitation, the induced level of humoral immunity directed toward the antigen is at least four-fold, and preferably at least 16-fold greater than the levels of the humoral immunity directed toward the antigen prior to the administration of the expressible polynucleotide of this invention to the subject. The immune response may also be measured qualitatively, by means of a suitable *in vitro* assay or *in vivo*, wherein an arrest in progression or a remission of neoplastic or infectious disease in the subject is considered to indicate the induction of a therapeutic immune response.

A further aspect of the invention is the administration of an expressible polynucleotide in combination with other forms of treatment including surgery, radiation therapy, and chemotherapy, to provide a vaccination against recurrence of cancer. By priming the immune system of subjects to recognize the cancer cells of their own cancers, the immune system will be more prepared to counter a subsequent regrowth, thus improving the prognosis for the subjects. This same type of vaccination can be used to improve a subject's resistance to recurrent diseases, including many parasitic and viral diseases. Thus, the expressible polynucleotides of the invention may be administered to stimulate a prophylactic immune response. The term "prophylactic immune response", as used herein, refers to an increase in long term humoral and/or cellular immunity, as measured by standard techniques, which is directed toward an antigen associated with the subject's disease condition. In general, the prophylactic immune response is measured qualitatively, wherein a delay in the onset of recurring system is achieved.

The subject treated in the method of the invention may be a human or non-human subject.

In various embodiments of the invention, the polynucleotide may encode a heat shock protein derived from the same species or alternatively from a different species relative to the species of the subject.

- 12 -

The cells into which the expressible polynucleotide are introduced may be neoplastic cells or other cancer cells from the subject to be treated or vaccinated. In particular, the method of the invention is usefully applied to treat or vaccinate subjects suffering from cancer including solid tumors and neoplastic disease such as sarcoma, lymphoma, carcinoma, leukemia and melanoma. The method of the invention may also be utilized to stimulate an immune response to infectious diseases, including parasitic, fungal, yeast, bacterial, mycoplasmal and viral diseases, where a particular class of cells can be identified as harboring the infective entity. For example, but not by way of limitation, the cells treated may be infected with a human papilloma virus, a herpes virus such as herpes simplex or herpes zoster, a retrovirus such as human immunodeficiency virus 1 or 2, a hepatitis virus, an influenza virus, a rhinovirus, respiratory syncytial virus, cytomegalovirus, adenovirus, *Mycoplasma pneumoniae*, a bacterium of the genus *Salmonella*, *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Clostridium*, *Escherichia*, *Klebsiella*, *Vibrio*, *Mycobacterium*, amoeba, a malarial parasite, *Trypanosoma cruzi*, etc. Thus, for example, in the case of human papilloma virus (HPV), the expressible polynucleotide could be introduced into epithelial cells infected with HPV from the subject to be treated.

The method of the invention can be practiced using various compositions. One such composition of matter is a recombinant vector comprising a promoter system effective to promote expression of the vector in mammalian cells; and a region encoding a XDEL-negative heat shock protein. The vector may be one in which the bases encoding a XDEL-signal have been deleted from the region encoding the heat shock protein, or one in which additional bases have been added to the region encoding the heat shock protein to mask the XDEL-signal in the expressed product. An exemplary vector would include gp96, preferably human gp96, as the heat shock protein.

- 13 -

A further composition of matter in accordance with the invention is a recombinant vector comprising a promoter system effective to promote expression of the vector in mammalian cells; and a region encoding a cytosolic heat shock protein such as hsp70 to which a signal comprising a positively-charged amino acid or group of a few amino acids at the amino terminal end of the peptide, a hydrophobic core region and a third region of greater polarity than the hydrophobic region is attached to promote uptake of the expressed cytosolic heat shock protein by the endoplasmic reticulum has been added.

The invention will now be further described by way of the following non-limiting examples.

EXAMPLE 1

PREPARATION OF KDEL DELETION MUTANTS

The preparation of mammalian expression vectors encoding full-length and KDEL-deleted Drosophila BiP cDNAs (dBiP, also referred to a grp78) is described in Munro et al., *Cell* 48: 899-907 (1987). The expression vector encoding dBiP with a KDEL deletion (KDEL-deleted dBiP, SAGM2) has the KDEL signal replaced with a *c-myc* sequence in a modified AHP2 plasmid. A further construct, WT dBiP (SAGMK1) has the *c-myc* sequence inserted between the KDEL signal and the balance of the dBiP.

Wild type mouse BiP cDNA was cloned into plasmid pBSBiP. Haas, *Proc. Nat'l Acad. Sci. USA* 85: 2250-2254 (1988). In this plasmid, the BiP sequence is flanked by BamH I sites. The BamH I fragment was therefore recovered from pBSBiP and cloned into the BamH I site of mammalian expression vector pCDNA3 (In-Vitrogen). Mouse KDEL-deleted BiP was constructed using pCDNA3 by cloning the Bgl II-EcoR V PCR fragment (nucleotide 1360 to nucleotide 1981) of BiP using two oligonucleotide primers complementary to BiP. The nucleotides coding for the last four amino acids (KDEL) were omitted, and a stop codon and an EcoR V restriction site put in its place.

- 14 -

EXAMPLE 2PREPARATION OF TUMOR CELLS EXPRESSING BiP

CMS-5 sarcoma cells, a methylocholanthrene-induced fibrosarcoma of BALB/c mouse origin, were adapted to culture and grown in DMEM medium (Gibco Life Technologies, Inc.) supplemented with 10% FCS. Sub-confluent monolayers (2×10^9 cells) were transfected with 2 ug of mammalian expression vector containing BiP or KDEL-deleted BiP using lipofectamine (Gibco) according to the manufacturer's directions. Briefly, the expression vector and 6 ul of lipofectamine were diluted separately in 100 uL serum-free medium (OPTI-MEM® I Reduced Serum Medium, Gibco-BRL). The two solutions were then mixed and incubated at room temperature for 45 minutes to allow formation of DNA-liposome complexes. 800 uL OPTI-MEM® was added to the complexes, mixed, and overlaid onto rinsed cells. After a 6 hour incubation at 37°C, 1 mL growth medium containing 20% FCS was added. Fresh medium was added to the cells 24 hours post-transfection.

Stable clones were selected by adding 800 ug/mL geneticin (Gibco-BRL) to the cells 72 hours later. The selection medium was changed every 3 days. Colonies of stably transfected cells were seen after 10 to 14 days. Expression of BiP for each clone picked was assayed for by radiolabeling. Newly synthesized BiP was detectable by immunoprecipitation and gel electrophoresis.

Acutely (transiently) BiP expressing clones were trypsinized and used for animal injection 60 hours post-transfection.

EXAMPLE 3VACCINATION WITH BiP EXPRESSING TUMOR CELLS

Freshly prepared acutely transfected CMS-5 sarcoma cells prepared in accordance with Example 2 were prepared for use in vaccination of mice by trypsinization and then washing three times in PBS. 2×10^6 cells were injected intradermally into the abdomen of CB6F-1/J [Balb/c x C57BL/6 F1] mice (Jackson Laboratories) and monitored for tumor growth over a 90 day period. The results are summarized in

- 15 -

Table 1 and shown graphically in Figs. 1A-H. As shown, injection with the parental CMS-5 sarcoma cells, with mock transfected cells, and with cells transfected with KDEL-deleted vectors led in all cases to the formation of tumors.

5 Tumors also developed in one control mouse where the cells were transfected with the vector only (no BiP). On the other hand, only one of six mice transfected with the complete wild-type drosophila BiP (including the *c-myc*) or wild-type mouse BiP developed tumors.

10

TABLE 1		
Cell Line	Number of Tumor Free Mice per number of mice injected	Percent Tumor Free Survivals
Parental CMS5	0/4	0%
Mock Transfected	0/3	0%
15 Vector Only	1/3	33%
Mouse BiP wild type	2/3	66%
Mouse BiP KDEL deleted	0/3	0%
Drosophila BiP wild type	3/3	100%
20 Drosophila BiP KDEL deleted	0/3	0%

20

EXAMPLE 4

The experiment of Example 3 was repeated using larger groups of mice and using two different clones of stable mouse BiP-expressing transfected cells. As shown in Table 2 and Figs 2A-F, the results were essentially the same, with none of the mice injected with the vector expressing wild-type mouse BiP having any detected tumor formation.

25

TABLE 2

Cell Lines	Number of Tumor Free mice per number mice injected	Percent Tumor Free Survivals
Parental CMS5	0/4	0%
Vector Only	4/15	27%
5 Mouse BiP wild type: clone 1	6/6	100%
Mouse BiP wild type: clone 2	11/11	100%
Mouse BiP KDEL deleted: clone 1	1/9	11%
Mouse BiP KDEL deleted: clone 2	2/15	13%

EXAMPLE 5

Tumor-free mice that had been injected with the wild-type BiP expression vector in the experiment of Example 4 were rechallenged after an interval of two months by injection of 2×10^6 or 10×10^6 CMS-5 sarcoma cells into the left flank of each mouse. Unimmunized CB6F-1/J mice of the same age group that had not been previously immunized were injected at the same time with 2×10^6 CMS-5 sarcoma cells as controls. As shown in Table 3, none of the previously immunized mice developed detectable tumors, while all three of the control mice developed tumors and died.

TABLE 3

Rejected Tumor	Number of Tumor Free Mice/Number of Mice Injected	% Survivals
25 Mouse BiP wild type: clone 1	3/3 3/3	100% 100%
Mouse BiP wild type: clone 2	5/5 6/6	100% 100%
Control	0/3	0%

- 17 -

EXAMPLE 6CONSTRUCTION OF VECTOR EXPRESSING gp96

To create an expressible vector encoding gp96, the coding region of murine GP96 (mGP96) was digested with BamH I, blunted with Klenow and cloned into the mammalian expression vector pRC/CMV (In-Vitrogen) which has been digested with Xba I and also blunted with Klenow. The resulting plasmid, p96/CMV was then used as template for generating a fragment spanning nucleotides 1653 to 2482 that lacks the nucleotides encoding the amino acids KDEL. The primers used for amplification of this fragment were

5' primer:

GCGGATCCTAGTTTAGACCAGTATGTC

[SEQ ID No. 1]

3' primer:

15 CCGAATTCTGGGCCCCAATTTACTCTGTAGATTCCTTTTCTGTTT [SEQ ID No. 2]

The restrictions sites for BamH I in the 5'-primer and EcoR I and Sal I in the 3'-primer are shown in bold italics. A stop codon (shown underlined) is provided four nucleotides from the Sal I restriction site in the 3'-primer. The BamH I/EcoR I PCR fragment was subcloned into pBlueScript II (Stratagene) to further amplify the amount of DNA. The amplified DNA was then digested with PflM I and Apa I, gel purified and subcloned into p96/CMV to give pMS215 which codes for KDEL-deleted gp96.

25

EXAMPLE 7VACCINATION WITH GP96 EXPRESSING VECTORS

Plasmids encoding the wt mGp96 (pMS216) and KDEL deleted mGp96 (215) are transfected into CMS-5 sarcoma cells as described in Example 2. 86 h post transfection, the acutely transfected CMS-5 sarcoma cells are trypsinized, washed in PBS and prepared for use in vaccination of mice as described in Example 3.

Various publications are cited herein, the contents of which are hereby incorporated by reference in their entireties.

35

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Sloan-Kettering Institute for Cancer Research

ROTHMAN, James E.
HARTL, F. Ulrich
HOE, Mee H.
HOUGHTON, Alan
TAKEUCHI, Yoshizumi
MAYHEW, Mark

(ii) TITLE OF INVENTION: Method for Treatment of Cancer and Infectious Diseases and Compositions Useful in Same

(iii) NUMBER OF SEQUENCES: 2

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Oppedahl & Larson
(B) STREET: 1992 Commerce Street Suite 309
(C) CITY: Yorktown
(D) STATE: NY
(E) COUNTRY: US
(F) ZIP: 10598

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette - 3.5 inch, 1.44 Mb storage
(B) COMPUTER: IBM compatible
(C) OPERATING SYSTEM: MS DOS
(D) SOFTWARE: Word Perfect

(vi) CURRENT APPLICATION DATA :

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION :

(A) NAME: Larson, Marina T.
(B) REGISTRATION NUMBER: 32,038
(C) REFERENCE/DOCKET NUMBER: MSK.P-018-WO

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (914) 245-3252
(B) TELEFAX: (914) 962-4330
(C) TELEX:

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: no

(iv) ANTI-SENSE: yes

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: mouse

(ix) feature

- 19 -

- (D) OTHER INFORMATION: amplification primer for gp96
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCGGATCCTA GTTTAGACCA GTATGTC. 27

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 44

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: no

(iv) ANTI-SENSE: no

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: mouse

(ix) feature

(D) OTHER INFORMATION: amplification primer for gp96

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCGAATTCGG GCCCAATTT ACTCTGTAGA TTCCTTTTCT GTTT 44

- 20 -

Claims

- 1 1. A method for stimulating a therapeutic immune
2 response in a subject in need of such treatment, comprising
3 the step of introducing to the subject, or to a target group
4 of cells from the subject, an expressible polynucleotide
5 encoding a eukaryotic heat shock protein.
- 1 2. A method for stimulating a prophylactic
2 immune response against recurrence of a disease from which a
3 subject is suffering, comprising administering to the
4 subject, or to a target group of cells from the subject
5 harboring an antigen associated with the disease, an
6 expressible polynucleotide encoding a eukaryotic heat shock
7 protein.
- 1 3. The method according to claim 1 or 2, wherein
2 the eukaryotic heat shock protein is an XDEL-negative heat
3 shock protein.
- 1 4. The method according to claim 1 or 2, wherein
2 the eukaryotic heat shock protein is a heat shock protein
3 from which the carboxy-terminal XDEL signal has been
4 deleted.
- 1 5. The method according to claim 1 or 2, wherein
2 the eukaryotic heat shock protein is a heat shock protein in
3 which the carboxy-terminal XDEL signal has been masked.
- 1 6. The method according to claim 1 or 2, wherein
2 the expressible polynucleotide encodes a eukaryotic heat
3 shock protein having a carboxy-terminal signal of the
4 sequence XDEL.
- 1 7. The method according to claim 6, wherein the
2 eukaryotic heat shock protein is BiP.
- 1 8. The method according to any of claims 1 to 7
2 wherein the eukaryotic heat shock protein is gp96.

- 21 -

1 9. The method according to any of claims 1 to 8,
2 wherein the cells are cancer cells.

1 10. The method according to any of claims 1 to 8,
2 wherein the cells are neoplastic cells.

1 11. The method according to claim 10, wherein the
2 neoplastic cells are selected from among sarcoma cells,
3 lymphoma cells, leukemia cells, carcinoma cells and melanoma
4 cells.

1 12. The method according to any of claims 1 to 8,
2 wherein the cells are infected with a virus.

1 13. The method according to any of claims 1 to 8,
2 wherein the cells are infected with a parasite.

1 14. The method according to any of claims 1 to 8,
2 wherein the cells are infected with a mycoplasma.

1 15. The method according to any of claims 1 to 8,
2 wherein the cells are infected with a bacterium.

1 16. The method according to any of claims 1 to 8,
2 wherein the cells are infected with a fungus or yeast.





1 17. The method according to any of claim 1 to 16,
2 wherein the expressible polynucleotide is introduced into a
3 target group of cells *ex vivo*, and the target group of cells
4 are thereafter administered to the subject.

1 18. The method according to claim 17, wherein the
2 composition is administered by transfection in a liposome.

1 19. The method according to any of claim 1 to 18,
2 wherein the subject is a human.

- 22 -

- 1 20. A recombinant vector comprising
2 (a) a promoter system effective to promote
3 expression of the vector in mammalian cells; and
4 (b) a region encoding a cytosolic heat shock
5 protein to which an amino terminal signal sequence
6 effective to promote uptake of the expressed cytosolic heat
7 shock protein by the endoplasmic reticulum has been added.
- 1 21. The recombinant vector of claim 20, wherein
2 the signal sequence comprises a positively charged N-
3 terminal region, a hydrophobic core region and a third
4 region of greater polarity than the hydrophobic region.
- 1 22. The recombinant vector according to claim 20
2 or 21, wherein the cytosolic heat shock protein is hsp70.
- 1 23. A method for stimulating an immune response
2 to an antigen present in a target group of mammalian cells,
3 comprising administering to the target group of cells an
4 expressible polynucleotide, wherein expression of the
5 polynucleotide is effective to increase the amount of
6 antigen-associated heat shock protein present in the cells.
- 1 24. The method according to claim 23, wherein the
2 expressible polynucleotide encodes a eukaryotic heat shock
3 protein.
- 1 25. The method according to claim 23 or 24,
2 wherein the expressible polynucleotide encodes a peptide
3 having a signal sequence recognized by the erd-2 receptors.
- 1 26. The method according to claim 25, wherein the
2 signal sequence is XDEL.






Fig. 1G




Fig. 1H




Fig. 1E




Fig. 1F

3/5




Fig. 2A




Fig. 2B

4/5

Clone 1 mouse BiP wild type




Fig. 2C

Clone 2 mouse BiP wild type




Fig. 2D

5/5

Clone 1 mouse BIP KDEL deleted




Fig. 2E

Clone 2 mouse BIP KDEL deleted




Fig. 2F

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13360

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 48/00; C12N 15/63
US CL :514/44; 424/93.21; 435/320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44; 424/93.21; 435/320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS, MEDLINE, EMBASE, BIOSIS, CAPLUS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94/11513 A1 (MEDICAL RESEARCH COUNCIL) 26 May 1994, see entire document.	1-7, 20-26
Y	LOWRIE et al. Towards a DNA vaccine against tuberculosis. Vaccine. 1994, Vol. 12, No. 16, pages 1537-1540, see entire document.	1-7, 20-26
Y	SILVA et al. A single mycobacterial protein (hsp 65) expressed by a transgenic antigen-presenting cell vaccinates mice against tuberculosis. Immunology. 1994, Vol. 82, pages 244-248, see entire document.	1-7, 20-26
Y	LUKACS et al. Tumor Cells Transfected with a Bacterial Heat-Shock Gene Lose Tumorigenicity and Induce Protection against Tumors. J. Exp. Med. July 1993, Vol. 178, pages 343-348, see entire document.	1-7, 20-26

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search
30 SEPTEMBER 1996

Date of mailing of the international search report
29 OCT 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

D. CURTIS HOGUE, JR.

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13360

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LOWRIE et al. Mycobacterium leprae HSP65 vaccinates mice against tuberculosis when expressed from the cloned gene in transplanted bone marrow cells. J. Cellular Biochemistry. 1995, Suppl. 19B, page 220, abstract No. B6-316, see entire document.	1-7, 20-26
A, P	EDGINGTON, S.M. Therapeutic Applications of Heat Shock Proteins. Bio/Technology. December 1995, Vol.13, pages 1442-1444, see entire document.	1-7, 20-26
Y, P	BLACHERE et al. Heat shock protein-based cancer vaccines and related thoughts on immunogenicity of human tumors. Seminars in Cancer Biology. 1995, Vol. 6, pages 349-355, see entire document.	1-7, 20-26

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13360

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 8-19
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.